

CO

11F

A study of the chemical composition of various divisions of the nervous system. III. The peripheral nerves of various divisions of the nervous system of the cow. A. V. Palladin, E. Ya. Raabbe and R. M. Gel'man. *Ukrain. Biokhim. Zhur.* 9, 160 (in Russian 1938; in English 1939-40); cf. *C. A.* 30, 5277. —The chem. compns. of the anterior and posterior roots of the spinal cord per unit of dry matter are essentially the same; per unit of wet matter cholesterol, lecithin and cephalin contents of the anterior roots are lower than that of the posterior. In comparison with other divisions of the nervous system the abn. cholesterol contents are much greater in both cases. The inverse is the case for lipoids, nitrogen, creatine, P and water contents in the peripheral nerve. The N and creatine contents of the sympathetic nerve are higher than those of other nerves. It is concluded that a high protein and a medium cholesterol content are characteristic for the nonmedullated nerves. The ganglia of the posterior roots have approx. the same compn. as all the others, with the exception of a higher cholesterol and unsatd. phosphatides. Phylogenetically the cholesterol and unsatd. phosphatides are higher in older than in younger divisions of the peripheral and of the central nervous systems; the reverse is true of water and protein contents. E. B. Stefanowsky


ASIN-114: METALLURGICAL LITERATURE CLASSIFICATION


PREGNANCY AND LACTATION


Arginine, arginase and creatine in the embryo tissues of vertebrates. A. V. Palladin and K. Ya. Raspbi-Ukra. Biokhim. Zhur. 10, 183-226 (in Russian 223-31, in English 234-42) (1937).—The arginine and creatine contents of the tissues of chick embryos diminish gradually with the progress of the embryonic development; min. values were found toward the moment of hatching; on the 40-50 days of postnatal development, they approach those characteristic for adults. The activity of arginase in the tissues of chick embryos and rabbits is highest in the early days of development and decreases gradually to the low values characteristic for adults. The arginase of embryo tissues is, in most cases, activated by salts of bivalent Mn; when its activity is low, it is always activated. The coexistence of large quantities of arginine and of a high activity of arginase suggests that the latter possibly takes part in the synthesis of arginine-creatine, proteins during tissue growth. The similarity of arginine and creatine curves and the slight formation of urea in the chick embryo suggest that creatine is formed from arginine. The activity of arginase is the same in the liver of chick embryos as in adult chickens. In the liver of rabbit embryos, it is very high. The change in the water content of the embryo liver is somewhat different from that of other embryo tissues. The min. quantity is noted on the 5th or 6th day after birth, increasing afterward. In rabbit

embryos, the water content in the liver on the 30th day of gestation is the same as in adult rabbits. Elllischer's hypothesis on the special significance of arginine exchange for tissue growth is supported by the very large arginine content and the high activity of arginase in embryo tissues during the early days of development. U. S.

APPENDIX D: DETAILED CATALOGUE LITERATURE CLASSIFICATION








Brain creatine during the embryonic development of vertebrates. A. V. Palladin and E. Ya. Hashiba. *J. Physiol. (U. S. S. R.)* 26, 203-76 (in English, 270) (1959). The content of creatine (1), total N and H₂O in the brains of rabbit and guinea-pig embryos is higher during the 2nd half of embryonic development than in mature animals. During development these values decrease until they reach those characteristic of mature animals during the 1st month of post-embryonic growth. The content of these substances in the hemispheres, cerebellum and medulla oblongata of cow embryos from the 3rd to the 9th month of growth is highest during the earlier stages, and reaches the maturity level in the 6th or 7th month of embryogenesis. Beginning with the 3rd month I is highest in the cerebellum, while the medulla oblongata is richest in total N and H₂O.

DETAILED POLITICAL LITERATURE CLASSIFICATION

APPROVED FOR RELEASE: Tuesday, August 01, 2000 CIA-RDP86-00513R0013442



RELEASED AND DECLASSIFIED 1985

Glycogen and phosphorus compounds in the muscles of chick embryos. B. Ya. Ruskin. Biokhim. J. (Ukraine) 13, 373-85 (in Russian; 883-75 in English, 587-9) (1969); cf. C. A. 63, 50009, 83801.—To det. the origin of lactic acid in the chick embryo muscles, at work during hatching, the relative amounts of glycogen in the liver and muscles were investigated. The resulting data did not furnish the answer. The glycogen on the 14th and 15th day of chick embryo development is 1-2%, rising on the 17th to 20th day, falling sharply on the 21st, remaining at this level, or below, for 2-3 days after hatching, when the chicks feed mainly on their yolk, and then begins to rise again. There is no particular increase of glycogen in the neck and leg muscles to account for the sharp rise of lactic acid in these muscles. It is possible that there is a rapid supply of glycogen by the liver, which becomes glycogen-poor at this period, or by local resynthesis. The quantity of P is 300-500 mg. per 100 g. of dry tissue. As in adult birds, the creatine P is very small, but the pyrophosphoric fraction is greater than in 3-4-month-old and adult chicks.
B. Gutov


AIAA SEA METALLURGICAL LITERATURE CLASSIFICATION

PROCESS AND PROPERTIES INDEX

Data on the carbohydrate metabolism in the muscles of chick embryos. E. Ya. Raskin. *Biochem. J. (Ukraine)* 13, 561-600 (in Russian, 1907-8; in English, 604-6 (1929).—Data collected from a series of detns. indicate that the source of a considerable quantity of lactic acid may be glucose, more intensely glycolyzed by the enzyme systems of the embryo working muscle, and present in the latter in a greater quantity than in the less intensely working muscle. Since the glycogen consumption *in vivo* is considerably less than that of glucose, R. is speculating on the manner of disappearance of glycogen both in anaerobic and in aerobic glycolysis of embryo muscles and also on the possible consumption of glycogen without the direct formation of lactic acid. B. Gutof

ASME-SEA METALLURGICAL LITERATURE CLASSIFICATION

APPROVED FOR RELEASE: Tuesday, August 01, 2000 CIA-RDP86-00513R0013442



PROCESS AND PROPERTIES

Glycolytic processes and phosphorylation of carbohydrates in muscles of rabbits and chicks during embryonic development. R. Ya. Nasilina. *Nauk. J. Ukrainsk. 15*, No. 2-3, 273-305 (in Russian, 262-4; in English, 294-5) (1940); cf. *C. A.* 34, 41201. --The embryo muscles split off inorg. P in anaerobic autolysis (*in vivo*). This was not observed to take place on adding 0.05 M NaF; P disappearing from the soln.; but there was an accumulation of the little-hydrolyzed intermediary P fractions, including lactate monophosphate (I), greatest on adding glycogen, remaining unchanged or dropping with glucose. The non-inorganic-phosph. fraction contained, in addn. to I and phosphoglyceric acids, some unidentified P compds., which usually increased after the autolysis. The abs. amt. of I in the embryo muscles was higher than in adults. The rabbit muscles gave the greatest lactic acid formation with glucose, but glycogen also reacted. The decrease in lactic acid with NaF was not investigated; this is ascribed to its dehydration into pyruvic acid. R. rejects the theory that the absence of phosphorolysis is due to a deficiency of the enzyme system, since the embryo muscles glycolyze and phosphorylate glycogen very well; but the intermediate products, typical for the usually fermenting glycogen, were absent during the glucose glycolysis. References. B. Gutoff

11F

AIA-LLA METALLURGICAL LITERATURE CLASSIFICATION

E-37-13442

SEARCHED

SERIALIZED

INDEXED

FILED

100000 74

SEARCHED

SERIALIZED

INDEXED

FILED

SEARCHED

SERIALIZED

INDEXED

FILED

RYZHKOV, P.Ya., inzh.; SHERSHEVSKAYA, R.M., inzh.; RASHBA, T.S., inzh.
Hardening spare metallurgical equipment parts at the Petrovskii plant.
Met. i gornorud. prom. no.3:76-80 My-Je '63. (MIRA 17:1)

RASHBA, Ye. Ya. and GOTOVSEVA, Ye. P.

Rashba, Ye. Ya. and Gotovtseva, Ye. P. - "The products of hydrolytic decomposition of polysaccharides in the brain", Ukr. biokhim. zhurnal, 1949, No. 1, p. 56-72, (In Ukrainian, resume in Russian), - Bibliog: 6 items.

SO: U-4630, 16 Sept. 53, (latopis 'Zhurnal nykh Statey, No. 23, 1949).

RASHBA, O. Ya.

Chemical Abst.
Vol. 48 No.8
Apr. 25, 1964
Biological Chemistry

✓ Products of polysaccharide hydrolysis in the brain.
O. Ya. Rashba and O. P. Cotoriseva (Inst. Biochem.,
Acad. Sci. Ukr. S.S.R., Kiev). Ukrainsk. Biokhim. Zhur.,
21, 61-66 (in Russian, 70-2)(1949).—Brain amylase in pure
form splits 60-70% of starch, resulting in dextrins of high-
and low-mol. wt. and maltose. Glycogen hydrolysis re-
sulted in similar polysaccharides. By means of pure culture
yeast fermentation, the presence of glucose and maltose
was demonstrated. Studies of perfused cow, rabbit, and dog
brains indicated the presence of active maltase. Expts.
with brain suspensions showed that maltase activity at
first proceeds at an increasing rate, then subsides, and finally
disappears. At the height of its activity it is accompanied
by lactic acid formation. Changes in P compds. run
parallel to accumulation of P-esters in the presence of mal-
tose. The rate of maltose glycolysis is conditioned by
activity of maltase. D. S. Levine

RACHEL, Ye. Ya.

34142. Prevrashcheniye dekstrinov v tkani mazga. Ukr. biokhim. Zhurnal, 1949, № 3, s. 247-56-. Na Ukr. yaz.-Rezyume na rus. yaz.- Bibliogr 16 nazv.

SU: Knizhnaya Letopis' № 6, 1955

RASHBA, YE.YA.

USSR (600)

Nervous System

Problems of histochemical investigation of glycogen of the normal nervous system.
Identification of biological properties and differences of neurons. A. L. Shabash.
Reviewed by Ye. Ya. Rashba, Ukr. biokhim. zhur., 22, No 1, 1950.

Monthly List of Russian Accessions. Library of Congress, October 1952. UNCLASSIFIED

RASHBA, E. Ya.

"Amylase and Its Role in the Initial Stages of Carbohydrate Metabolism in the Cerebrum of Certain Animals." Sub 27 Apr 51, Acad Med Sci USSR.

Dissertations presented for science and engineering degrees in Moscow during 1951.

SO: Sum. No. 480, 9 May 55

RASHIN, S.YA.

Isolation and study of the chemical composition of nuclei
of the cerebral tissue. A.V. Falleldin, S.IA. Rashin,
TS.M. Shtutman. Ukr. biokhim. zhur. 23 no. 3:265-277 '51.

PALLADIN, O.V.; RASHBA, Ye.Ya. [reviewers]; PENFIELD, W.; RASMUSSEN, Th. [authors].

"Cerebral cortex of man. A clinical study of localization of function."
W.Penfield, Th.Rasmussen. Reviewed by O.V.Palladin, I.E.IA.Rashba. Ukr.
biokhim.zhur. 24 no.2:258-260 '52. (MLRA 6:11)
(Brain)

RASEBA, O.Ya.

Role of nucleic acids in variability of microorganism. Mikrobiol.
zhur. 15 no.1:59-65 '53. (MIRA 7:3)

1. Z Institutu mikrobiologii AN URSR.
(Bacteria) (Nucleic acids) (Variation (Biology))

RASHBA, O.Ya.

"Bacteriochemistry." E.M.Gubarev. Reviewed by O.IA.Rashba. Ukr.biokhim.
zhur. 25 no.2:237-241 '53. (MLR 6:6)
(Biochemistry) (Bacteria)

USSR.

Changes in the nucleic acids of artificially induced variants of *Escherichia coli*. O. Ya. Rastin and T. O. Galkina. *Mikrobiol. Zhur., Akad. Nauk Ukr. R.S.R.*, 16, No. 1, 27-32 (Russian summary, 32-3)(1964).—*E. coli* 103 and *Salmonella breslau* 353 were used. The nucleic acids of both original cultures were determined. *E. coli* variants were obtained by growing the original 103 strain in a culture medium containing *S. breslau* autolyzate (cf. Yizir, et al., *Ibid.*, 15, No. 3 and 4 (1953)). Variants 125, 125M (obtained by passing 125 through a mouse), 38, 106, and 60 were almost identical with *S. breslau* in serological, immunological, radio-fermentation, and pathogenic properties. Cells of these variants were washed and total N and nucleic acids were determined by the method of Schmidt and Thannhauser (*C.J.B.*, 40, 231). Results are expressed in terms of % of ribonucleic (I) and deoxyribonucleic (II) acid P per mg. of bacterial cells. *E. coli* 103 and *S. breslau* 353 contained about the same amount of I but II was higher in *S. breslau*. Variants 125 and 106 contained less of I and more of II than either *E. coli* 103 or *S. breslau* 353. Content of I and II in variants 125 M and 60 was about same as in *E. coli* 103. The content of nucleic acids in variant 38 was not constant, but the amt. of II approximated that in *S. breslau*. Each expt. was repeated 10 times. B. S. Levin.

KASHIN D. J.

M.S.R.

Deaminase of purine and of ribonucleic acid in *Escherichia coli* variants grown in the presence of *Salmonella breslau* autolysates. O. Ya. Rashba and L. S. Krachko (Inst. Microbiol., Acad. Sci. Ukr. S.S.R.). Mikrobiol. Zhur. Akad. Nauk Ukr. R.S.R. 16, No. 1, 34-40 (Russian summary, 39-40) (1954). *S. breslau* 353 and *E. coli* 163 and its variants 38, 106, 125, and 125 M were used. Suspensions of 24 hr. resting cultures were freed from substrate, and adenine (I), glutamine (II), or ribonucleic acid (III) added at the rate of 3-5 mg. per vol. of bacterial suspension contg. 8-8 mg. of bacterial N; pH was kept at 7-8. Suspensions of resting bacteria were incubated at 37° for 2 hrs. Deaminase activity was detd. from the NH₃ split off. Following the incubation period, the proteins were pptd. with trichloroacetic acid, and NH₃ in the centrifugate was detd. by isothermic distn. Split off NH₃ was found in the substrate-free suspension of all bacteria, the amt. detd. and subtracted correspondingly from the NH₃ detd. in the substrate suspnsion. During the 2-hr. incubation, deaminase of *E. coli* 163 and of *S. breslau* 353 split off 0.018-0.040% NH₃ N, or 5-16% of the N of amino groups of I, and 0.080-0.137 mg. or 50-81% of the N of amino groups of III. No deamination of II was detected. Variant 125 gave similar results. Variants 38 and 106 showed the presence of a slight guanase activity; otherwise their deaminase activity was as above. Variant 125 M differed sharply from its parent culture *E. coli* 163 and from *S. breslau* 353. Its guanase activity was high (0.140-0.334 mg. NH₃ N, or 40-85%) and the deaminase for I was 5-10 times as high as that of *E. coli* 163 or *S. breslau* 353.

B. S. I.

USSR

The biochemical properties of the antibacterial substances of some labiates. O. V. Bushkin, S. I. Zekpuikha, T. P. Mandrik, and M. B. Kagan'ska. Mikrobiol. Zhur., Akad. Nauk Ukr. R.S.R., 16, No. 2, 62-6 (Russian summary, 60 X 1954).—*Thymus serpyllum* var. *potosicus* (I), *Teucrium chamaedrys* (II), and *Stachys recta* (III), used in popular and homeopathic therapy, were investigated for their antibacterial properties in fresh and dried state. The usual chemophytological methods were employed to obtain the following fractions: ether exts., acidic substances, basic substances, saponins and glucosides, tannins. Exts. suspected of having tannic acid were tested for di- and trinhydroxyphenolic hydroxy groups. For the antibacterial tests *Micrococcus pyogenes* var. *aureus* 209, the Hiss-Flexner bacillus, and *Microbacterium* 5 were used. I contained at least 2 substances with antibacterial properties: the ether exts., which arrested the growth of *M. pyogenes* var. *aureus* 209 in 10^{-4} - 25×10^{-5} dilns., and a low mol. tannic substance isolated from the isoec. sediment of the boiling H₂O ext. of the plant, which arrested the growth of *M. pyogenes* var. *aureus* 209 at 10^{-1} - 15×10^{-2} dilns. and the Hiss-Flexner bacillus at 25×10^{-1} . II also had 2 antibacterial substances: an ether ext. obtained during the period of blooming (ether ext. of plant during nonblooming period) contained no antibacterial substances which arrested the growth of *M. pyogenes* var. *aureus* 209 only, and a low mol. wt. tannin-type of substance, which contained procathechins and was acidic. III contained low mol. wt. acidic substances with antibacterial properties belonging to the group of tannophenolic acids.

B. S. Levine

VASHI
USSR.

The biochemical properties of the antibacterial substances of *Inula britannica* and of *Solidago virgaurea*. O. Ya. Rashba, S. I. Zelepukha, and M. B. Kagan's'ka. Mikrobiologiya, Akad. Nauk Ukr. R.S.R. 16, No. 2, 67-8 (1964) (Russian summary, 69).—Both plants are popularly used in wound and ulcer healing. The plants were dried and extd. The fatty and other ether fractions of the roots and above-ground parts of *I. britannica* had antibacterial properties, but not at all times. Similar exts. of *S. virgaurea* were more potent against all test microorganisms. The alc. ext. of the isolect. sediment (of hot H₂O ext. acidified) exhibited similar antibacterial properties. B. S. Levine

PHS/BA, 07

The nucleoproteins of *Salmonella breslau*. Some of their properties and methods for their isolation. O. Ya. Busha, Mikrobiol. Zhur., Akad. Nauk Ukr. R.S.R. 18, No. 3, 43-7 (Russian summary, 48)(1954).—Fresh and desiccated masses of cells of *Salmonella breslau* 353 were extd. with saline and alk. solns., the exts. combined, and the nucleoproteins pptd. by acidifying to the isoelec. point. Ppts. and mother liquors were studied. The nucleoprotein can be extd. with 0.1M NaHCO₃, 1.4M NaCl (ribonucleoproteins), 0.010-0.012M NaOH, and 0.1M Na citrate (mixed nucleoproteins). An increase in the concn. of the NaOH soin. increases the gross fraction extd. but the nucleoprotein fraction remains the same. Nucleoproteins extd. from dry bacteria and pptd. with trichloroacetic acid become relatively insol.; they can be redissolved in 0.5% (0.05M) soin. of NaOH. Ground frozen masses of *S. breslau* extd. with 0.1M Na citrate become polymerized. Bacteria washed with physiol. saline yielded a nucleoprotein which contained less deoxyribonucleic acid than was obtained from similar extn. of whole bacteria. The fractionation of the nucleic acids by the method of Belozerskil and Chernomordnikova (cf. C.A. 35, 14579) yielded poor results, but gave higher deoxyribonucleic acid values. The cytoplasmic protein sepd. from the nucleoproteins proved to be ribonucleoproteins. The nucleoproteins here dealt with can be adsorbed on asbestos filter disks. B. S. Levine

RASHBA, O.Ya.

✓ 8000. Alterations of nucleic acid content of colon bacillus on assimilation of Breslau bacillus complete antigen and nucleoprotide. O. Ya. Rashba and T. O. Caffine. Mikrobiol. zh., 1955, 21, 17, 14-17; Zh. mikrobiol., 1956, Abstr. No. 75816. Regardless of the character and intensity of the changes of bacteriological, serological and pathological properties of variants obtained, their content of nucleic acid differed little from that of the original culture. Variants obtained on media with complete antigen and nucleoprotide of Breslau bacillus, differed less from the original cultures and were less diverse in character, than variants obtained on media with autolysates of Breslau bacillus. (Ukrainian, Russian Summary) *C. PAULCE*

Inst. Microbiol. AS Ukr SSR

RASHBA, O. Ya.

Methods for separation of deoxyribonucleic acid from
Bacteria Breslau. O. Ya. Rashba. *Mikrobiol. Zhur. Akad.*
Nauk Ukr. S.S.R., 18, No. 1, 47-50 (in Ukrainian). Summary
summary, 61 (1953).—Attempts to sep. deoxyribonucleic
acid (I) from Bacteria Breslau 553 (*Salmonella typhimurium*)
gave the following results. The modified method of Mc-
Carty and Avery (C.A. 40, 2194) gave very little pure I
and it was not polymeric. Treatment of the bacterial mass
with alkali at high temp. gave partly dephosphorylated I.
Autolysis of the bacteria in presence of citrate did not
give reproducible results. All preps. of I obtained were
toxic for *Escherichia coli*.
A. Sementsov

Rashba, O.YA.

The activity of deaminase of purines and of nucleic acids
in different *Escherichia coli* and *Salmonella typhimurium*.
O. Ya. Rashba and S. M. Tsiuskalovska. *Mikrobiol.
Zhurn. Akad. Nauk Ukr. S.S.R., Inst. Mikrobiol. im. D.K.
Zabolotnogo* 18, No. 2, 30-4 (Russian summary, 38) (1958). —
A study was made of the deaminase of adenine, guanine,
ribonucleic, and deoxyribonucleic acids of 6 strains of *E. coli*
and 6 strains of *S. typhimurium*. These strains were of dif-
ferent origins isolated at different times between 1946 and
1952. All strains were typical culturally, serologically, and
morphologically. The bacteria were grown on beef-pep-
tone-agar, harvested and washed. For the detn. of de-
aminase activity a sample of the washed bacteria was
added to a substrate mixt. contg. adenine hydrochloride,
guanine, and ribonucleic acid and incubated for 2 hrs.;
the inc contg. deoxyribonucleic acid was incubated for 24
hrs. The amt. of NH₃ split off was detd. Control tests
were also run. *E. coli* strains SK and sr. deaminated ade-
nine at a rate 2-3 times below that of other strains. During
the period of 1952-3 the activity of adenase and guanase
had considerably increased. All strains of *E. coli* deaminate
ribonucleic acid at nearly the same rate. Deoxyribonucleic
acid is deaminated at a slow but regular rate by strain 163
only. *S. typhimurium* deaminates the purines at slower
rate than most *E. coli* strains and at approx. the same
rate as strains SK and sr. *S. typhimurium* deaminates ribo-
nucleic acid at a rate similar to *E. coli*. Deoxyribonucleic
acid is deaminated by *S. typhimurium* faster than by *E.
coli*.

B. S. Levine

2

Med

RASHES A, O. YA.

Med ✓ The respiration rate and fermentation intensity of bacterial variants developed in the filtrate of *Salmonella typhimurium*. O. Ya. Rushin. *Mikrobiol. Zhur., Akad. Nauk Ukr. S.S.R., Inst. Mikrobiol. im D. K. Zabolotnogo* 18, No. 3, 30-5 (Russian summary, 36) (1960). — The respiration rate and fermentation intensity of the variants were compared with those of the parent cultures. Cultures were grown on beef-peptone-agar for 18 hrs. at 37°, and further treated as described by Cook and Stephenson (C.A. 23, 1661). The bacterial suspension contained 30×10^9 cells/ml. Respiration and fermentation rates were detd. by the Warburg method. Glucose (0.05M) was added to some of the tests. Data were recorded in terms of ml.^{-1} of gas consumed or eliminated/hr./0.1 g. of bacterial N (Q_{O_2} and Q_{CO_2}). On the basis of the results obtained in relation to the rate of respiration and the intensity of fermentation it was possible to place the variants into 4 groups: (1) 4 cultures differing very slightly from the parent cultures; (2) 7 cultures in which the respiration and fermentation rates, although clearly discernible, were at levels below those of the parent cultures; (3) 2 cultures whose fermentation intensity exceeded that of the parent cultures; (4) 5 cultures which did not utilize glucose either for respiration or fermentation, but metabolized amino acids and some other substrates. The processes of respiration and of fermentation of the variants differed only slightly from those of the parent *S. typhimurium*. Cells which were unable to utilize glucose manifested a notable endogenous respiration and were able to utilize a no. of other substrates. Some cultures had slow growth rates, and neutralized developing acidity by their ability to break down N-contg. compds. of the medium.

B. S. Levine

Inst. Microbiology AS Ukr SSR

Rashba, O. Ya.

The utilization of phosphoric acid by *Streptococcus lactis* grown in milk. O. Ya. Rashba and S. I. Novikova. *Mikrobiol. Zhur., Akad. Nauk Ukr. S.S.R., Inst. Mikrobiol. im. D. K. Zabolotnogo* 18, No. 3, 37-8 (Russian summary, 39) (1950); cf. *C.A.* 51, 2935c.—A study was made of the content of inorg. phosphoric acid in the fraction of easily hydrolyzable phosphate esters and of the reducing substances in sterile milk inoculated with intensive acid producers and weak acid producers. From the data obtained the content of lactose was calc'd. Simultaneously, as the lactose of the milk is being consumed by *S. lactis*, the inorg. phosphoric acid content is being reduced, and the fraction of the easily hydrolyzable phosphate compds. is increased. Towards the end of the expt., when the milk becomes thick, the quantity of inorg. phosphoric acids returns to normal. It is assumed that these changes are a reflection of the fact that the phosphoric acids are being drawn into the process of lactose breakdown by the proliferating culture of *S. lactis*. B. S. Levine

Rashba, O. Ya.

Distr: 4E4c/4E3d

Diffusion of current carriers in a semiconductor in the presence of an external electric field. O. Ya. Rashba. Soviet Phys. Tech. Phys. 1, 1381-7(1957) (English translation); Zhur. Tekh. Fiz. 26, 1415-18(1958). -- Exact formulas are derived for determining the distribution of nonequilibrium holes (photoholes) in a slab and a plate of semiconductors in the presence of a homogeneous elec. field. Illumination is carried out by point or filament light probes for a momentary and a stationary source. Solutions are obtained for the case of simultaneous presence of vol. and surface recombination. E. E. Bouvier.

21

2
2

JRC

Rashba, Ye.Y.
DAVIDOV, O.S.; RASHBA, Ye.Y.

Light absorption in molecular crystals on weak interaction between excitons and phonons [in Ukrainian with summaries in Russian and English]. Ukr.fiz.zhur. 2 no.3:226-241 J1-S '57. (MIRA 10:10)

1. Institut fiziki AN URSR.
(Excitons) (Crystals--Optical properties)

BASEBA, O.Ya.

Academician Aleksei Nikolaevich Bach. Mikrobiol.shur. 19 no.1:
59-62 '57. (MLRA 10:?)
(BACH, ALEKSEI NIKOLAEVICH, 1857-1946)

KASHBA, O. Ye.

Studying phytocides of *Helichrysum arenarium*. Microbiol. zhur. 19
no.2:25-28 '57. (KRA 10:3)

1. Z Institutu mikrobiologii AN URSR.

(PLANTS

phytocides of *Helichrysum arenarium*, antibact. eff.)

(ANTIBACTERIAL

phytocides of *Helichrysum arenarium*)

RASHBA, I.C. Ya.

EXCERPTA MEDICA Sec 4 Vol 12/3 Med. Micro. Mar 59

825. A STUDY OF NUCLEOPROTEINS OF B. COLI VARIANTS OBTAINED BY ASSIMILATION OF B. BRESLAU PRODUCTS (Russian text) - Rashba H. J. and Kaganskaya M. B. Dept. of Microbe Biochem., Inst. of Microbiol., Acad. of Scis of the Ukrainian SSR, Kiev - BIOKHIMIYA 1957, 22/6 (1008-1012) Graphs 1 Tables 1

A study was carried out of the nucleoproteins of 3 E. coli variants obtained on media containing S. breslau autolysates. The nucleoproteins of E. coli and S. breslau differ in viscosity and in isoelectric precipitation curves. The nucleoproteins of the variants of E. coli had the same low viscosity as the S. breslau nucleoproteins, and a similar shape of the curve of isoelectric precipitation, but with some larger additional maximum in the acid zone.

RASHBA, Ye.Ya.

Certain aspects of metabolism in cultures not requiring glucose
isolated from filtrates of Bacterium breslau [with summary in English]
Mikrobiologija, 26 no.3:277-284 My-Je '57. (MIRA 10:10)

1. Institut mikrobiologii AN USSR, Kiyev.
(*SALMONELLA*, metabolism,
breslau, metab. in strains not requiring glucose (Eus))

RASHBA, Ye. Ya.

25-2-13/43

AUTHORS: Bel'tyukova, K.I. and Rashba Ye.Ya., Scientific Workers of the Institute of Microbiology, AS Ukrainian SSR

TITLE: Useful Properties of Everlasting (Poleznyye svoystva bessmertnika)

PERIODICAL: Nauka i Zhizn', 1958, # 2, pp 45-46 (USSR)

ABSTRACT: A group of co-workers of the Institute of Microbiology of the Ukrainian SSR Academy of Sciences, engaged in research on Ukrainian flora, disclosed that everlasting grown on sands contains two substances having antibacterial properties. One of them is the color of the plant - helichrysine; the other is a resinous substance called arenarine, which is of great importance in controlling the bacteria of agricultural plants. Arenarine, for instance, proved to be very effective in destroying tomato cancer without being detrimental to the tomato plant itself. Also, it exerts a favorable influence on the germination of tomato seeds. In all cases, whether the tomato plants were sprinkled with a solution of arenarine (1:10,000) or the seeds were treated with this antibiotic before seeding, the yields proved to be

Card 1/2

Useful Properties of Everlasting

25-2-13/43

by 11-26% higher than without applying arenarine.

ASSOCIATION: Institute mikrobiologii AN USSR (Institute of Microbiology,
AS Ukrainian SSR)

AVAILABLE: Library of Congress

Card 2/2

RASHBA, Ye.Ya. [Rashba, O.IA]

Oxidation of glucose and alanine by bacteria of the intestinal group producing an alkaline reaction. Mikrobiol.zhur. 20 no.3:
(MIRA 11:11)
49-54 '58

1. Iz instituta mikrobiologii AN USSR.
(GLUCOSE)
(ALANINE)
(ENTEROBACTERIACEAE)
(ALCALIGENES FAECALIS)

RASHKA, Ye.Ya. [Rashba, O.IA.]

Problems and future development of the biochemistry of micro-
organisms. Mikrobiol.zhur. 21 no.1:17-20 '59. (MIRA 12:6)
(MICROORGANISMS, metabolism,
research in Russia (Uk))

RASHBA, Ye.Ya. [Rushba, O.IA.]

Modern data on the role of nucleic acids in the variability and
heredity of micro-organisms. Mikrobiol.zhur. 21 no.1:61-67
'59. (MIRA 12:6)

(NUCLEIC ACIDS, metab.

microorganisms, eff. on variability & hered..
review (Uk))

(MICROORGANISMS, metab.

nucleic acids, eff. on variability & hered..
review (Uk))

RASHBA, Ye.Ya. [Rashba, O.IA.]

Enzymes utilized in the initial stages of carbohydrate metabolism
by *Bacterium alkalescens*. *Mikrobiol.zhur.* 21 no.2:57-61 '59.
(MIRA 12:9)

(*SHIGELLA* - metabolism)
(*ENZYMOSS* - metabolism)

KASHTBA, X. Ya.

SOV/30-59-1-51
Soviet Union
Afrimyan, M. K. et al. Candidates of Biological Sciences

Title: Use of Antibiotics in Plant Cultivation (Primeneniye antibiotikov v rasstoyedovedenii).

Periodical: Vestnuk Akademii Nauk SSSR, 1959, Ser. 1, pp 142-143 (USSR)

Abstract: A conference dealing with this subject took place in Yerevan from 5 to 10 October 1958. It had been called by the Institute of Microbiology, All-Union Agricultural Research Institute of the Academy of Sciences of the USSR (USSR), the All-Union Selectivity and Breeding Research Institute, the Veterinary Institute for Agricultural Microbiology of the USSR, and the Sakharovskiy Agricultural Academy and Research Inst. (Institute for Microbiology of the Academy of Sciences of the Armenian SSR).

M. K. Afrimyan spoke about antibiotic metabolites which promote the development of higher plants.

I. V. Pashkevich reported on investigations of several factors which produce active antibiotics against the disease of potato tuber rot and diplodia in potato.

A. G. Karapetyan spoke about the utilization of antibiotics in fighting potato ring rot and various diseases of vegetables.

G. G. Bubumyan spoke on the effect of preparations from culture of streptomycetes to prevent wilt of the cotton bush.

A. G. Tsvetkov, I. V. Lopatin, N. A. Bokhanova, Ya. S. Mikhaleva, and G. G. Bubumyan spoke about the production of antibiotic preparations which produce active antibiotics against diseases of vegetable cultures and plants with a short life-span.

N. P. Chikaturova, T. A. Gerasimova, and G. G. Bubumyan dealt with the utilization of epiphyte microflora in fighting several fungi diseases in plants.

D. G. Chikaturova, N. P. Chikaturova, I. Z. Shatschikova, and G. G. Bubumyan developed remedies obtained in investigation of phytopathogenicity as well as its utilization in fighting diseases concerning the cotton bush and beans.

B. P. Gulyashyan, T. P. Petrosyan, L. G. Arakelyan, and G. G. Bubumyan spoke on the effect of antibiotic preparations of various types against bacterial cankers in fighting diseases of decorative plants.

T. G. Gulyashyan, G. J. Melikyan, described the investigation of plant antibiotics.

S. A. Karapetyan spoke about the production of the preparation "Primenilin" and "Mithobolin" and their effectiveness against various diseases in wheat, oats and rye.

A. G. Karapetyan reported on results obtained in the utilization of antibiotic bacteria against oilseed.

Card 2/4

Card 3/4

Card 4/4

Card 5/4

Card 6/4

Card 7/4

Card 8/4

Card 9/4

Card 10/4

Card 11/4

Card 12/4

Card 13/4

Card 14/4

Card 15/4

Card 16/4

Card 17/4

Card 18/4

Card 19/4

Card 20/4

Card 21/4

Card 22/4

Card 23/4

Card 24/4

Card 25/4

Card 26/4

Card 27/4

Card 28/4

Card 29/4

Card 30/4

Card 31/4

Card 32/4

Card 33/4

Card 34/4

Card 35/4

Card 36/4

Card 37/4

Card 38/4

Card 39/4

Card 40/4

Card 41/4

Card 42/4

Card 43/4

Card 44/4

Card 45/4

Card 46/4

Card 47/4

Card 48/4

Card 49/4

Card 50/4

Card 51/4

Card 52/4

Card 53/4

Card 54/4

Card 55/4

Card 56/4

Card 57/4

Card 58/4

Card 59/4

Card 60/4

Card 61/4

Card 62/4

Card 63/4

Card 64/4

Card 65/4

Card 66/4

Card 67/4

Card 68/4

Card 69/4

Card 70/4

Card 71/4

Card 72/4

Card 73/4

Card 74/4

Card 75/4

Card 76/4

Card 77/4

Card 78/4

Card 79/4

Card 80/4

Card 81/4

Card 82/4

Card 83/4

Card 84/4

Card 85/4

Card 86/4

Card 87/4

Card 88/4

Card 89/4

Card 90/4

Card 91/4

Card 92/4

Card 93/4

Card 94/4

Card 95/4

Card 96/4

Card 97/4

Card 98/4

Card 99/4

Card 100/4

Card 101/4

Card 102/4

Card 103/4

Card 104/4

Card 105/4

Card 106/4

Card 107/4

Card 108/4

Card 109/4

Card 110/4

Card 111/4

Card 112/4

Card 113/4

Card 114/4

Card 115/4

Card 116/4

Card 117/4

Card 118/4

Card 119/4

Card 120/4

Card 121/4

Card 122/4

Card 123/4

Card 124/4

Card 125/4

Card 126/4

Card 127/4

Card 128/4

Card 129/4

Card 130/4

Card 131/4

Card 132/4

Card 133/4

Card 134/4

Card 135/4

Card 136/4

Card 137/4

Card 138/4

Card 139/4

Card 140/4

Card 141/4

Card 142/4

Card 143/4

Card 144/4

Card 145/4

Card 146/4

Card 147/4

Card 148/4

Card 149/4

Card 150/4

Card 151/4

Card 152/4

Card 153/4

Card 154/4

Card 155/4

Card 156/4

Card 157/4

Card 158/4

Card 159/4

Card 160/4

Card 161/4

Card 162/4

Card 163/4

Card 164/4

Card 165/4

Card 166/4

Card 167/4

Card 168/4

Card 169/4

Card 170/4

Card 171/4

Card 172/4

Card 173/4

Card 174/4

Card 175/4

Card 176/4

Card 177/4

Card 178/4

Card 179/4

Card 180/4

Card 181/4

Card 182/4

Card 183/4

Card 184/4

Card 185/4

Card 186/4

Card 187/4

Card 188/4

Card 189/4

Card 190/4

Card 191/4

Card 192/4

Card 193/4

Card 194/4

Card 195/4

Card 196/4

Card 197/4

Card 198/4

Card 199/4

Card 200/4

Card 201/4

Card 202/4

Card 203/4

Card 204/4

Card 205/4

Card 206/4

Card 207/4

Card 208/4

Card 209/4

Card 210/4

Card 211/4

Card 212/4

Card 213/4

Card 214/4

Card 215/4

Card 216/4

Card 217/4

Card 218/4

Card 219/4

Card 220/4

Card 221/4

Card 222/4

Card 223/4

Card 224/4

Card 225/4

Card 226/4

Card 227/4

Card 228/4

Card 229/4

Card 230/4

Card 231/4

Card 232/4

Card 233/4

Card 234/4

Card 235/4

Card 236/4

Card 237/4

Card 238/4

Card 239/4

Card 240/4

Card 241/4

Card 242/4

Card 243/4

Card 244/4

Card 245/4

Card 246/4

Card 247/4

Card 248/4

Card 249/4

Card 250/4

Card 251/4

Card 252/4

Card 253/4

Card 254/4

Card 255/4

Card 256/4

Card 257/4

Card 258/4

Card 259/4

Card 260/4

Card 261/4

Card 262/4

Card 263/4

Card 264/4

Card 265/4

Card 266/4

Card 267/4

Card 268/4

Card 269/4

Card 270/4

Card 271/4

Card 272/4

Card 273/4

Card 274/4

Card 275/4

Card 276/4

Card 277/4

Card 278/4

Card 279/4

Card 280/4

Card 281/4

Card 282/4

Card 283/4

Card 284/4

Card 285/4

Card 286/4

Card 287/4

Card 288/4

Card 289/4

Card 290/4

Card 291/4

Card 292/4

Card 293/4

Card 294/4

Card 295/4

Card 296/4

Card 297/4

Card 298/4

Card 299/4

Card 300/4

Card 301/4

Card 302/4

Card 303/4

Card 304/4

Card 305/4

Card 306/4

Card 307/4

Card 308/4

Card 309/4

Card 310/4

Card 311/4

Card 312/4

Card 313/4

Card 314/4

Card 315/4

Card 316/4

Card 317/4

Card 318/4

Card 319/4

Card 320/4

Card 321/4

Card 322/4

Card 323/4

Card 324/4

Card 325/4

Card 326/4

Card 327/4

Card 328/4

Card 329/4

Card 330/4

Card 331/4

Card 332/4

Card 333/4

Card 334/4

Card 335/4

Card 336/4

Card 337/4

Card 338/4

Card 339/4

Card 340/4

Card 341/4

Card 342/4

Card 343/4

Card 344/4

Card 345/4

Card 346/4

Card 347/4

Card 348/4

Card 349/4

Card 350/4

Card 351/4

Card 352/4

Card 353/4

Card 354/4

Card 355/4

Card 356/4

Card 357/4

Card 358/4

Card 359/4

Card 360/4

Card 361/4

Card 362/4

Card 363/4

Card 364/4

Card 365/4

Card 366/4

Card 367/4

Card 368/4

Card 369/4

Card 370/4

Card 371/4

Card 372/4

Card 373/4

Card 374/4

Card 375/4

Card 376/4

Card 377/4

Card 378/4

Card 379/4

Card 380/4

Card 381/4

Card 382/4

Card 383/4

Card 384/4

Card 385/4

Card 386/4

Card 387/4

Card 388/4

Card 389/4

Card 390/4

Card 391/4

Card 392/4

Card 393/4

Card 394/4

Card 395/4

Card 396/4

Card 397/4

Card 398/4

Card 399/4

Card 400/4

Card 401/4

Card 402/4

Card 403/4

Card 404/4

Card 405/4

Card 406/4

Card 407/4

Card 408/4

Card 409/4

Card 410/4

Card 411/4

Card 412/4

Card 413/4

Card 414/4

Card 415/4

Card 416/4

Card 417/4

Card 418/4

Card 419/4

Card 420/4

Card 421/4

Card 422/4

Card 423/4

Card 424/4

Card 425/4

Card 426/4

Card 427/4

Card 428/4

Card 429/4

Card 430/4

Card 431/4

Card 432/4

Card 433/4

Card 434/4

Card 435/4

Card 436/4

Card 437/4

Card 438/4

Card 439/4

Card 440/4

Card 441/4

Card 442/4

Card 443/4

Card 444/4

Card 445/4

Card 446/4

Card 447/4

Card 448/4

Card 449/4

Card 450/4

Card 451/4

Card 452/4

Card 453/4

Card 454/4

Card 455/4

Card 456/4

Card 457/4

Card 458/4

Card 459/4

Card 460/4

Card 461/4

Card 462/4

Card 463/4

Card 464/4

Card 465/4

Card 466/4

Card 467/4

Card 468/4

Card 469/4

Card 470/4

Card 471/4

Card 472/4

Card 473/4

Card 474/4

Card 475/4

Card 476/4

Card 477/4

Card 478/4

Card 479/4

Card 480/4

Card 481/4

Card 482/4

Card 483/4

Card 484/4

Card 485/4

Card 486/4

Card 487/4

Card 488/4

Card 489/4

Card 490/4

Card 491/4

Card 492/4

Card 493/4

Card 494/4

Card 495/4

Card 496/4

Card 497/4

Card 498/4

Card 499/4

Card 500/4

Card 501/4

Card 502/4

Card 503/4

Card 504/4

Card 505/4

Card 506/4

Card 507/4

Card 508/4

Card 509/4

Card 510/4

Card 511/4

Card 512/4

Card 513/4

Card 514/4

Card 515/4

Card 516/4

Card 517/4

Card 518/4

Card 519/4

Card 520/4

Card 521/4

Card 522/4

Card 523/4

Card 524/4

Card 525/4

Card 526/4

Card 527/4

Card 528/4

Card 529/4

Card 530/4

Card 531/4

Card 532/4

Card 533/4

Card 534/4

Card 535/4

Card 536/4

Card 537/4

Card 538/4

Card 539/4

Card 540/4

Card 541/4

Card 542/4

Card 543/4

Card 544/4

Card 545/4

Card 546/4

Card 547/4

Card 548/4

Card 549/4

Card 550/4

Card 551/4

Card 552/4

Card 553/4

Card 554/4

Card 555/4

Card 556/4

Card 557/4

Card 558/4

Card 559/4

Card 560/4

Card 561/4

Card 562/4

Card 563/4

Card 564/4

Card 565/4

Card 566/4

Card 567/4

Card 568/4

Card 569/4

Card 570/4

Card 571/4

Card 572/4

Card 573/4

Card 574/4

Card 575/4

Card 576/4

Card 577/4

Card 578/4

Card 579/4

Card 580/4

Card 581/4

Card 582/4

Card 583/4

Card 584/4

Card 585/4

Card 586/4

Card 587/4

Card 588/4

Card 589/4

Card 590/4

Card 591/4

Card 592/4

Card 593/4

Card 594/4

Card 595/4

Card 596/4

Card 597/4

Card 598/4

Card 599/4

Card 600/4

Card 601/4

Card 602/4

Card 603/4

Card 604/4

Card 605/4

Card 606/4

Card 607/4

Card 608/4

Card 609/4

Card 610/4

Card 611/4

Card 612/4

Card 613/4

Card 614/4

Card 615/4

Card 616/4

Card 617/4

Card 618/4

Card 619/4

Card 620/4

Card 621/4

Card 622/4

Card 623/4

Card 624/4

Card 625/4

Card 626/4

Card 627/4

Card 628/4

Card 629/4

Card 630/4

Card 631/4

Card 632/4

Card 633/4

Card 634/4

Card 635/4

Card 636/4

Card 637/4

Card 638/4

Card 639/4

Card 640/4

Card 641/4

Card 642/4

Card 643/4

Card 644/4

Card 645/4

Card 646/4

Card 647/4

Card 648/4

Card 649/4

Card 650/4

Card 651/4

Card 652/4

Card 653/4

Card 654/4

Card 655/4

Card 656/4

Card 657/4

Card 658/4

Card 659/4

Card 660/4

Card 661/4

Card 662/4

Card 663/4

Card 664/4

Card 665/4

Card 666/4

Card 667/4

Card 668/4

Card 669/4

Card 670/4

Card 671/4

Card 672/4

Card 673/4

Card 674/4

Card 675/4

Card 676/4

Card 677/4

Card 678/4

Card 679/4

Card 680/4

Card 681/4

Card 682/4

Card 683/4

Card 684/4

Card 685/4

Card 686/4

Card 687/4

Card 688/4

Card 689/4

Card 690/4

Card 691/4

Card 692/4

Card 693/4

Card 694/4

Card 695/4

Card 696/4

Card 697/4

Card 698/4

Card 699/4

Card 700/4

Card 701/4

Card 702/4

Card 703/4

Card 704/4

Card 705/4

Card 706/4

Card 707/4

Card 708/4

Card 709/4

Card 710/4

Card 711/4

Card 712/4

Card 713/4

Card 714/4

Card 715/4

Card 716/4

Card 717/4

Card 718/4

Card 719/4

Card 720/4

Card 721/4

Card 722/4

Card 723/4

Card 724/4

Card 725/4

Card 726/4

Card 727/4

Card 728/4

Card 729/4

Card 730/4

Card 731/4

Card 732/4

Card 733/4

Card 734/4

Card 735/4

Card 736/4

Card 737/4

Card 738/4

Card 739/4

Card 740/4

Card 741/4

Card 742/4

Card 743/4

Card 744/4

Card 745/4

Card 746/4

Card 747/4

Card 748/4

Card 749/4

Card 750/4

Card 751/4

Card 752/4

Card 753/4

Card 754/4

Card 755/4

Card 756/4

Card 757/4

Card 758/4

Card 759/4

Card 760/4

Card 761/4

Card 762/4

Card 763/4

Card 764/4

Card 765/4

Card 766/4

Card 767/4

Card 768/4

Card 769/4

Card 770/4

Card 771/4

Card 772/4

Card 773/4

Card 774/4

Card 775/4

Card 776/4

Card 777/4

Card 778/4

Card 779/4

Card 780/4

Card 781/4

Card 782/4

Card 783/4

Card 784/4

Card 785/4

Card 786/4

Card 787/4

Card 788/4

Card 789/4

Card 790/4

Card 791/4

Card 792/4

Card 793/4

Card 794/4

Card 795/4

Card 796/4

Card 797/4

Card 798/4

Card 799/4

Card 800/4

Card 801/4

Card 802/4

Card 803/4

Card 804/4

Card 805/4

Card 806/4

Card 807/4

Card 808/4

Card 809/4

Card 810/4

Card 811/4

Card 812/4

Card 813/4

Card 814/4

Card 815/4

Card 816/4

Card 817/4

Card 818/4

Card 819/4

Card 820/4

Card 821/4

Card 822/4

Card 823/4

Card 824/4

Card 825/4

Card 826/4

Card 827/4

Card 828/4

Card 829/4

Card 830/4

Card 831/4

Card 832/4

Card 833/4

Card 834/4

Card 835/4

Card 836/4

Card 837/4

Card 838/4

Card 839/4

Card 840/4

Card 841/4

Card 842/4

Card 843/4

Card 844/4

Card 845/4

Card 846/4

Card 847/4

Card 848/4

Card 849/4

Card 850/4

Card 851/4

Card 852/4

Card 853/4

Card 854/4

Card 855/4

Card 856/4

Card 857/4

Card 858/4

Card 859/4

Card 860/4

Card 861/4

Card 862

RASHBA, Ye.Ya. [Rashba, O.IA]; MOSTOVOVA, G.A. [Mostovova, H.O.]

Dependence of the antibacterial properties of arenarine on the
time of harvesting of immortelle and on other factors. Mikro-
biol.zhur. 24 no.2:48-55 '62. (MIRA 15:12)

1. Institut mikrobiologii AN UkrSSR.
(EVERLASTING FLOWERS) (MATERIA MEDICA, VEGETABLE)

RASHBA, Ye.Ya. [Rashba, O.IA.]

Some components of arenarin and their antibacterial effects.
Mikrobiol. zhur. 26 no.2:26-31 '64. (MIRA 18:8)

1. Institut mikrobiologii AN UkrSSR.

PASHBA, Ye.Ya. [Pashba, O.I.]

Effect of the chromatographic fractions of arenarin on shoots of
some plants and on *Corynebacterium michiganense*. Mikrobiol. zhur.
(MIRA 18:8)
27 no.4:61-66 '65.

1. Institut mikrobiologii i virusologii AN UkrSSR.

AYZENMAN, B.Ye. [Aizenman, B.IU.]; SHVAYGER, M.O.; MANDRIK, T.P.;
BREDIKHINA, A.N. [Bredikhina, A.M.]; ORISHCHUK, I.P. [Orishchuk, I.P.];
KOLESOVA, E.A. [Kolesova O.A.]; MISHENKOVA, Ye.L. [Mishenkova, O.L.];
GALKINA, T.A. [Halkina, T.O.]; ZAKHAROVA, I.Ya.; RASHBA, Ye.Ya.
[Rashba, O.IA.]; LAUSHNIK, G.M. [Laushnyk, H.M.];
PREOBRAZHENS'KA, N.Ye. [Preobrazhens'ka, N.IU.]

Effect of substances of bacterial origin on Ehrlich's carcinoma.
Mikrobiol. zhur. 27 no.6:61-67 '65. (MIRA 19:1)

1. Institut mikrobiologii i virusologii AN UkrSSR.

RASHBA, Ye.Ya.; KAGANSKAYA, M.B.

Study of the electrophoretic properties of citrate-soluble
proteins of the colon bacillus in its variations. Biokhimiia
30 no.1:3-6 Ja-F '65. (MIRA 18:6)

1. Institut mikrobiologii AN UkrSSR, Kiyev.

RASHEA, Ye.Ya. [Rashba, O.Ia.]; FAUZMIK, G.M. [Fauzmyk, H.M.]

Lipopoly saccharides of alkali-forming of Enterobacteriaceae.
Mikrobiol. zhur. 26 no.3:22-27 '64. (MIR 18:5)

1. Institut mikrobiologii AN UkrSSR.

RASHEVA, Ye.Ya. [Rashba, G.IA.]; KOLCHINSKAYA, I.B. [Kolchinsk'a, I.B.];
ZAKHAROVA, I.Ya.; MATYSHENSKAYA, M.S. [Matyshens'ka, M.S.]

First All-Union Biochemical Congress. Mikrobiol. zhur. 26
no.3:94-100 '64. (MIRA 18:5)

TYDEL'SKAYA, I.L.; MYSLAVSKAYA, I.S.; RASHBA, Ye.Ya.; ZAKHAROVA, I.Ya.

Study of C-precipitinogen in atypical streptococcal strains. Zhur.
mikrobiol., epid.i immun. 40 no.12:93-97 D '64.

(MIRA 17:12)

1. Iz Ukrainskogo instituta klinicheskoy meditsiny imeni Strazhesko
i Instituta mikrobiologii AN UkrSSR.

RASHBA, Ye.Ya [Rashba, S. IA]; ZAYCHIKOV, A.M. [Zaychikov, O.M.]

Study of the amylase and saccharase activity of mycelial wastes
from the penicillin and streptomycin. Mikrobiol. zhur. 24 no.6:
32-36 '62 (MIRA 17:5)

1. Institut mikrobiologii AN UkrSSR.

Kazan, Tatar. (Russia.)

Specific polysaccharides of streptococci. Mikrobiol. zhur. 25
no.11A-51 163. (MIRA 1785)

PIDOPLICHKO, Ivan Grigor'yevich[Pidoplichko, I.H.]; RASHBA, Yelena
Yakovlevna[Rashba, O.IA.]; KILEROG, N.M.[Klieron, N.M.],
red.

[Origin and development of life on the earth] Pokhodzhen-
nia i rozvytok zhyttia na zemli. Kyiv, Naukova dumka,
1964. 73 p. (MIRA 18:2)

BEL'TYUKOVA, Klavdiya Ignat'yevna; RASHBA, Yelena Yakovlevna;
KULIKOVSKAYA, Mariya Dmitriyevna; MATSHEVSKAYA, Mariya
Stepanovna; GVOZDYAK, Roatislav Il'ich; DROBOT'KO, V.G.,
akademik, otv. red.; SKUTSKAYA, M.P., -red.izd-va;
KADASHEVICH, O.A., tekhn. red.

[Arenarin and its use in plant growing] Arenarin i ego pri-
menenie v rastenievodstve. Kiev, Izd-vo AN USSR, 1963. 1
163 p. (MIRA 16:8)

(Arenarin) (Plant diseases)

RASHBA, Ye.Ya. [Rashba, D.IA.]

Fifth International Biochemistry Congress. Mikrobiol. zhur. 23 no.6:
70-71 '61. (MIRA 15:4)
(BIOCHEMISTRY--CONGRESSES)

RASHBA, Yelena Yakovlevna; VIZIR, P.Ye.[Vizyr, P.IE.], doktor biol.
nauk, otv. red.; SKUTSKA, N.P., red. izd-va; LISOVETS, O.M.
[Lysovets', O.M.], tekhn. red.

[Biology and biochemistry of the alcaligenic bacteria of the
intestinal group] Biologiya i biokhimia luhoutvoruiuchykh
bakterii kyshkovoi grupy. Kyiv, Vyd-vo Akad.nauk URSR, 1961. 130 p.
(MIRA 15:7)

(*Alcaligenes faecalis*)

ASHBA, YE. A. (USSR)

"Carbohydrate Exchange of Alkali-Forming Bacteria of the Intestinal Group."

Report presented at the 5th International Biochemistry Congress,
Moscow, 10-16 Aug 1961

RASHBA, Ye.Ya.; GALKINA, T.A.; ZAKHAROVA, I.Ya.; KAGANSKAYA, M.B.

Biochemical changes observed in certain coli bacteria during
variability. Trudy Inst. mikrobiol. no. 6:102-109 '59.
(MIRA 13:10)

1. Institut mikrobiologii AN USSR.
(SALMONELLA TYPHIMURIUM) (ESCHERICHIA COLI)

RASHBA, Ye.Ya. [Rashba, O.IA.]; LAUSHNIK, G.M. [Laushnyk, H.M.]

Decomposition of some phosphorus organic compounds by B.
megatherium var. phosphaticum.. Mikrobiol. Zhur. 22 no. 1:22-26
'60. (MIRA 13:10)

1. Institut mikrobiologii AN USSR.
(BACILLUS MEGATHERIUM) (PHOSPHORUS ORGANIC COMPOUNDS)

| | |
|------------|---|
| Country | : USSR |
| Category | : Microbiology. General Microbiology. Growth and Development of the Microbial Population. |
| Abs. Jour | : Ref Zhur-Biol., No 23, 1958, No 103602 |
| Author | : Rashba, Ye. Ya. |
| Institut. | : -- |
| Title | : Certain Features of Metabolism in Cultures which Do Not Consume Glucose Isolated from Filtrates of the Breslau Bacillus |
| Orig. Pub. | : Mikrobiologiya, 1957, No 2, 277-284 |
| Abstract | : The process of respiration and fermentation were studied manometrically in original Breslau bacillus strains and in regenerated cultures obtained from filtrates of them. It was established that the biochemically inert regenerated cultures which are incapable of using glucose oxidize pyruvate and, in some cases, sodium succinate and also oxidize and deaminate alanine, glycocoll, and asparagine. Therefore, the usual test of inoculating regenerated cultures onto "a variegated series" for the determination of their biochemical activity is inadequate. In the author's opinion, the lack of ability of these cultures to split sugar is associated with the absence or poor development of the enzyme systems in them which carry out the initial |
| Card: | 1/1 |

RASHBE, E.I.: SHEKA, V.I.

Combined resonance of band electrons in crystals with a
sphalerite lattice. Fiz. tver. tela 3 no.6:1735-1749
Je '61.

(MIRA 14:7)

1. Institut poluprovodnikov AN USSR, Kiyev.
(Electrons) (Sphalerite) (Crystal lattices)

RASHBA, E.I.; SHEKA, V.I.

Combined resonance in n-type InSb. Fiz. tver. tela 3
no.6:1863-1870 Je '61. (MIRA 14:7)

1. Institut poluprovodnikov AN USSR, Kiyev.
(Indium antimonide)

RASHCHENKO, I.D.; BALAKIN, F.S., dots.

Tissue regeneration after surgical treatment of cows with narrowed milking ducts. Veterinariia 36 no.2:80-81 F '59. (MIRA 12:2)

1. Glavnnyy veterinarnyy vrach Shalinskoy rayvetlechebnitsy Sverdlovskoy oblasti (for Rashchenko). 2. Sverdlovskiy gosudarstvennyy meditsinskiy institut (for Balakin).
(Udder--Diseases)

RASHCHENKO, I.D., veterinarnyy vrach.

Surgical method of eliminating the condition of hard milking
in cows. Veterinariia 32 no.8:68-70 Ag '55. (MLRA 8:10)

1.Ordona Lenina sovkhоз "Istek", Aramil'skogo rayona, Sverdlevskoy oblasti.
(UDDER--ABNORMALITIES AND DEFORMITIES)

RASHCHENKO, I. N.

Changes in the chemical constitution of the tubers of the Jerusalem artichoke in the process of sprouting. I. N. Raschenko, Trudy Kazakh. So'skotches. Issn. 9, No. 1, 73-85 (1955); Referat. Zhur., Khim., Biol. Khim. 1957, No. 5035.—A study was made of the changes in the mineral substances (I), carbohydrates (II) and proteins (III) of the tubers of *Helianthus tuberosus* in the different phases of its growth from the time of planting to the time of new tuber formation. In the process of growth the content of I in the tubers increased from 4.5% on the basis of dry wt. at the time of planting to 16.3% at the time of the stem development and to 32.4% at the time of the new tuber formation. At the corresponding periods of growth, II became sharply reduced from 78% to 45% and to 2%, resp. The content of III remained unchanged. Sucrose and monosaccharides constituted the basic parts of the II of the tubers; fructose predominated among the monosaccharides. In the process of growth the sucrose and the monosaccharides became quantitatively reduced.

B. S. Levine

RASHCHENKO, I.N.

Enriching the tubers of Jerusalem artichoke with yeast. Trudy Inst.
mikrobiol. i virus. Ak Kazakh SSR 2:88-94 '58 (MIRA 11:10)
(JERUSALEM ARTICHOKE)
(YEAST)

USSR/Farm Animals - General Problems.

Q-i

Abs Jour : Ref Thur - Biol., No 13, 1953, 33261

Author : Rashchenko, I.N.

Inst : Institute of Microbiology and Virology, AS KazSSR.

Title : Yeast-Fermentation of Girasole Tubers [*Jerusalem Artichokes*].

Orig Pub : Tr. In-ta mikrobiol. i virusol. AN KazSSR, 1958, 2, 33-94

Abstract : No abstract.

Card 1/1

- 2 -

RASHCHENKO, Ivan Nazarovich; SHERMAN, R.N., red.; OYSTRAKH, V.G.,
tekhn. red.

[Homemade cured food, preserves, and marinades] Domashnie
solen'ia, varen'ia i marinady. Alma-Ata, Kazakhskoe gos.izd-
vo, 1962. 221 p. (MIRA 16:2)

(Canning and preserving)

RASHCHENKO, Ivan Nazar'yevich; SHARIPOVA, N.G., red.; TURABAYEV, B., tekhn. red.

[Processing and preservation of vegetables, and fruits under home conditions] Pererabotka i khranenie ovoshchey, plodov v domashnikh usloviakh. Izd.2., dop. Alma-Ata, Kazgospizdat, 1963. 237 p. (MIRA 17:2)

RASHCHENKO, Ivan Mazarovich; GORDIYENKO, N.S., kand. sel'skokhozyaystvennykh
nauk, red.; LYAKHOVETSAYA, T.Ye., red.; KOZLOV, S.V., tekhn. red.

[Harvesting and ensiling corn] Uborka i silosovanie kulturasy. Pod
red. N.S. Gordienko. Alma-Ata, Kazakhskoe gos. izd-vo, 1956. 22 p.
(Kazakhstan--Corn (Maize)) (MIRA 11:7)
(Ensilage)

RASHCHENKO, N.I.

Secular reduction in the radio emission from Cassiopeia A. Priroda
50 no.7:97-98 Jl '61. (MIRA 14:6)

1. Gosudarstvennyy astronomicheskiy institut im. P.K.Shternberga,
Moskva. (Radio astronomy)

ROMADIN, G.; RASHCHEVSKO, N., ekonomist

State farms of Uzbekistan do not utilize hidden potentialities.
Fin.SSSR. 20 no.11:38-41 N '59. (MIRA 12:12)

1. Nachal'nik otdela Ministerstva finansov Uzbekskoy SSR
(for Romadin).
(Uzbekistan--State farms)

USSR / Forestry. Dendrology.

K-2

Abs Jour: Ref Zhur-Biol., No 6, 1958, 24884.

Author : Rashchenko, S. L.

Inst : Not given.

Title : Forestry of Kabardino-Balkariya.

Orig Pub: Uch. zap. Kabardino-Balkarsk n.-i. in-ta, 1957,
13, 81-109.

Abstract: No abstract.

Card 1/1

33

RASHCHEPKIN. A.P.

Field in the stator-to-rotor gap of an induction machine with
variable linear loading of the winding. Mag. Nizdr. No. 3:96-102
'65. (MIRA 18:30)

L 14230-66

ACC NR: AP5024908

UR/0382/65/000/003/0096/0102

AUTHOR: Rashchepkin, A.P.

ORG: None

TITLE: Gap field at a variable linear winding load of an induction machine

SOURCE: Magnitnaya gidrodinamika no. 3, 1965, 96-102

TOPIC TAGS: linear induction machine, induction motor design, winding theory, winding design optimization

ABSTRACT: In the stator-to-rotor air gap of a linear machine with an open magnetic flow circuit, the mission of the winding is to create a traveling magnetic field. However, the large gap present in the magnetic circuit creates additional pulsating fields in the working gap, causing deterioration of performance. The present paper analyses the spacial and time structure of the magnetic field in the working gap for the case of linear current density variation along the yoke. A compensating device with definite advantages over known compensators is proposed. It consists of 1) an ordinary three-phase winding with an optimized current amplitude, for which a design expression is given; and 2) a supplementary winding. The optimized design attains a current distribution resulting in an essentially pure traveling field in the working gap. Orig.art. has 2 figures, 17 formulas.

SUB CODE: 1309 SUBM DATE: 29Mar65/

ORIG REF: 006 OTH REF: 001

Card 1/1 QC

UDC 538.4:621.689

L 1003C-67 EWT(1)/EWP(m) WW
ACC NR: AP6034585 SOURCE CODE: UR/0382/66/000/003/0106/0110

AUTHOR: Rashchepkin, A. P.

ORG: none

TITLE: Longitudinal edge effect of induction machines

SOURCE: Magnitnaya gidrodinamika, no. 3, 1966, 106-110

TOPIC TAGS: edge effect, induction machine

ABSTRACT: The effect of a magnetic and electrical shunt of the terminal points of a linear induction machine has been investigated for one-dimensional flow of conductive fluid with a constant flow velocity. The conditions of perfect compensation of the longitudinal edge effect are found, and circuits are presented for improving such machines. The results obtained show the high electrical conductivity of the squirrel cage outside the active zone to be the basic cause of impairment of parameters of the induction machine equipped with the squirrel cage rotor. Orig. art. has: 2 figures and 14 formulas. [Based on author's abstract]

SUB CODE: 20/SUBM DATE: 29Jul65/ORIG REF: 008/

Card 1/1 egk

UDC: 621.313.39:538.4

ASATURYAN, A.Sh.; RASHCHEPKIN, K.Ye.; PETROVA, L.N.

Pipelines under stress. Izv. vys. ucheb. zav.; neft i gaz no.8:97-105
'58. (MIRA 11:10)

1.Moskovskiy neftyanoy institut im. akad. I.M. Gubkina i Bashkirskiy
nauchno-issledovatel'skiy institut neftyanoy promyshlennosti.
(Pipelines) (Strains and stresses)

RASHCHEPKIN, K.Ye.

Calculating the strained condition of a pipeline in major repairs.
Trudy NIITransneft' no.1:287-294 '61. (MIRA 16:5)
(Pipelines—Maintenance and repair) (Strains and stresses)

RASHCHEPKIN, K.Ye.; BARCHAN, N.I.; TIMERBAYEV, N.Sh.

Mechanized removal of protective coatings from pipelines. Trudy
NIITransneft' no.1:295-303 '61. (MIRA 16:5)
(Pipelines) (Protective coatings)

RASHCHEPKIN, K.Ye.; BARCHAN, N.I.

Analyzing the degree of the mechanization of labor consuming
processes in the major repair of pipelines. Trudy NIITransneft'
no.1:304-314 '61. (MIRA 16:5)
(Pipelines--Maintenance and repair)

RASHCHEPKIN, K.Ye.; BARCHAN, N.I.

Mechanized cleaning of pipelines in service during major repairs.
Neft.khoz. 39 no.1:54-58 1 Ja '61. (MIRA 17:3)

RASHCHEPAIN, K.Ye.; BEREZIN, V.L.

Stressed state of a working pipeline during overhauling.

Izv. vya. uch. zav.; neft' i gaz 5 no.9:77-81 '62.

(MIRA 17:5)

1. Ufimskiy neftyanyj institut.

YEDIGAROV, S.G.; LEVENTSOV, A.N.; KRUGLOV, A.N.; RASHCHEPKIN, Y.Ye.;
OVCHINNIKOV, I.S.

Mechanization of the packaging of solid petroleum bitumens.
Neft. khoz. 40 no.4:60-65 Ap '62. (MIRA 15:5)
(Bitumen)
(Packaging machinery)

YEDIGAROV, S.G.; RASHCHEPKIN, K.Ye.; OVCHINNIKOV, I.S.

Complete mechanization of major repairs of pipelines. Neft.
khoz. 40 no.10:55-62 0 '62. (MIRA 16:7)

(Pipelines—Maintenance and repair)

BEREZIN, V.L.; GUMEROV, A.G.; RASHCHEPKIN, K.Ye.

Performance of petroleum-plant tanks. Transp. i khran.
nefti no. 3:19-21 '63. (MIRA 17:7)

1. Ufimskiy neftyanoy institut i Nauchno-issledovatel'skiy
institut po transportu i khraneniyu nefti i nefteproduktov.

RASHCHEPKIN, K.Ye.; RAMEYEV, M.K.; VALIYEV, D.M.

Consolidated UIM-14 insulation machine. Transp. i khran. nefti
no.5:8-9 '63. (MIRA 17:3)

1. Nauchno-issledovatel'skiy institut po transportu i zhivotnyyu
nefti i nefteproduktov.

BEREZIN, V.L.; RASHCHEPKIN, K.Ye.

Stresses in operating pipelines occurring in major repairs
as a result of the deformation of tamped trenches. Izv. vys.
ucheb. zav.; neft' i gaz 6 no.4:71-74 '63. (MIRA 16:7)

1. Ufimskiy neftyanoy institut.
(Pipelines—Maintenance and repair)
(Strains and stresses) (Earthwork)

BEREZIN, V.I.; RASHCHEPKIN, N.Ye.; YASIN, E.M.

Calculation of stresses in the wall of a pipe in nonsymmetrical
pipeline hoisting. Izv. vys. ucheb. zav.; neft' i gaz 6 no.7:
95-101 '63. (MIRA 17:8)

1. Ufimskiy neftegazovoy institut.

BEREZIN, V.L.; RASHCHEPKIN, K.Ye.; YASIN, E.M.

Selection of boundary conditions in lifting an infinite
pipeline. Izv.vys.ucheb.zav.; neft' i gaz 6 no. 12:69-74
'63. (MIRA 17:5)

I. Ufimskiy neftyanoy institut.

YEDIGAROV, S.G.; RASHCHEPKIN, K.Ye.; MAYSKIY, A.A.

Mechanization of excavation in the major repair of pipelines. Transp.
i khran. nefti no.10:3-5 '63. (MIRA 17:9)

1. Nauchno-issledovatel'skiy institut po transportu i khraneniyu nefti
i nefteproduktov.

RASHCHEPKIN, K. Ye.; SULTANMURATOV, Kh. F.; TIMERBAYEV, N. Sh.; RAMEYEV,
M. K.

Investigating the operation of the vertical screw pumps of the
UIM-14 machine for applying protective coatings. Transp i khran
nefti no. 11:6-11 '63. (MIRA 17:5)

1. Nauchno-issledovatel'skiy institut po transportu i khraneniyu
nefti i nefteproduktov.

RASHCHEPKIN, K.Ye.; BUZHINSKIY, V.L.; OVCHINNIKOV, I.S.

OMS..1 combined high-speed cleaning device. Neft. khoz. 41
no.2:51-55 F '63. (MIRA 17:8)

L 31874-65 ENT(d)/ENT(m)/EMP(w)/EMA(d)/EMP(v)/EMP(k)/EHA(h) P-4/Peb EM
ACCESSION NR: AR5005871 S/0124/64/000/011/V012/V012

SOURCE: Ref. zh. Mekhanika, Abs. 11V95

AUTHORS: Yedigarov, S. G.; Rashchepkin, K. Ye.; Yasin, E. M.

TITLE: Stability of cylindrical shell under a load that is unevenly distributed along the generatrix

CITED SOURCE: Tr. N.-i. in-t po transp. i khraneniyu nefti i nefteproduktov, vyp. 3, 1961, 25-34

TOPIC TAGS: cylindrical shell, shell stability, thin shell

TRANSLATION: A solution is presented for the problem of stability in the small of an elastic thin circular cylindrical shell, loaded by a transverse pressure that is unevenly distributed over the length. The well-known stability equation of V. Z. Vlasov is used (Izv. AN SSSR. Otd. tekhn. n., 1947, No. 1, 27-52; Obshchaya teoriya obolochek i veye primeneniye v tekhnike [General Theory of Shells and Its Use in Engineering] M., Gosstekhizdat, 1949; Izbrannyye trudy [Selected Works] v. 1, M., Izd-vo AN SSSR, 1962, 364). The solution is obtained by the Bubnov-Galerkin

Card 1/2

L 31874-65

ACCESSION NR: AR5005871

method. The sag is chosen in the form

$$w = \sin np \sum_{m=1}^M w_m \sin \frac{mx}{l}$$

The load function is expanded in a Fourier series in cosines (with constant term). Expressions for the Fourier coefficients are given for three types of load. A calculation example is given. E. I. Grigolyuk.

SUB CODE: AS

ENCL: 00

Card 2/2

BEREZIN, V.L.; RASHCHEPKIN, K.Ye.; TIMURBAYEV, N.Sh.; YASIN, E.M.;
SULTANZIRATOV, Kh.F.; GUMEROV, A.G.; ZAKHAROV, I.Ya.

Experimental study of tension state of a pipeline during
capital repair. Izv. vys. ucheb. zav.; neft' i gaz 7 no.10:
89-91 '64. (MIRA 18:2)

1. Ufimskiy neftyanoy institut.

RASHCHEPKIN, K.Ye.; YASIN, E.M.

Stresses in a raised pipeline with fastened sections. Transp. i
khran. nefti i nefteprod. no.9;10-12 '64. (MIRA 17:10)

1. Nauchno-issledovatel'skiy institut po transportu i khraneniyu
nefti i nefteproduktov.

YEDIGAROV, S.G.; VOLZH, I.R.; RASHCHEPKIN, K.Ye.; MYSKII, A.A.;
VAL'YEV, B.M.; LEVKIVSKY, G.P.; ISMAGILEVA, P.B..

Excavator for uncovering pipelines in the ground. Transp. i khran.
nefti i nefteprod. no.10:12-14 '64.

(MIRA 17:12)

1. Nauchno-issledovatel'skiy institut po transportu i khraneniyu
nefti i nefteproduktov.

ZAKHAROV, I.Ya.; RASHCHEPKIN, K.Ye.

APPROVED FOR RELEASE: Tuesday, August 01, 2000 CIA-RDP86-00513R001344

Capital repairs of pipelines laid in swamps. Transp. i khran.
nefti i nefteprod. no.12:13-15 '64. (MIRA 18:2)

1. Nauchno-issledovatel'skiy institut po transportu i khraneniyu
nefti i nefteproduktov.

SULTANMURATOV, Kh.F.; RASHCHEPKIN, K.Ye.; YASIN, E.M.

Reducing pipeline hoist stresses. Transp. i khran. nefti i nefteprod.
no.8:5-9 '65. (MIRA 18:9)

1. Nauchno-issledovatel'skiy institut po transportu i khraneniyu nefti
i nefteproduktov.